

On Electromagnetic Production of Strange and Charm Mesons

Wednesday, 19 May 2021 18:05 (20 minutes)

New models for photo- and electroproduction of kaons on the proton were constructed [1,2,3] utilizing new experimental data from LEPS, GRAAL, and particularly CLAS collaborations. The higher spin nucleon (spin-3/2 and spin-5/2) and hyperon (spin-3/2) resonances were included using a consistent formalism and they were found to play an important role in the data description. In these analyses, we paid close attention to model predictions of the cross section at small kaon angles which are vital for accurate calculations of the hypernucleus-production cross section.

In order to account for the unitarity corrections at the tree level, we have introduced energy-dependent widths of nucleon resonances, which affect the choice of hadron form factors and the values of their cutoff parameters extracted in the fitting procedure.

We have implemented a new shape of electromagnetic form factors so that we are now able to describe also the process of electroproduction [4]. Moreover, for a reliable description of $K^+ \Lambda$ electroproduction at small Q^2 within our models it is necessary to take into account also a longitudinal coupling of virtual photons to nucleon resonances.

For the investigation of kaon photoproduction off the proton target, we have exploited also the hybrid Regge-plus-resonance (RPR) model [3] which provides an acceptable description of data in and above the resonance region. A novel feature of our version of the RPR model consists in applying a different scheme for the gauge-invariance restoration [5], which results in a need for a contact current. We reveal that the choice of the gauge-invariance restoration method may play a significant role for cross-section predictions at forward angles where data are scarce.

The sets of chosen nucleon resonances in our recent models are mutually quite well consistent and they also greatly overlap with the set selected in the Ghent analysis [6]. The results of our new isobar and RPR models will be compared with photo- and electroproduction experimental data and the properties of the models will be discussed.

The RPR approach was utilized to study also the \bar{D}^0 meson photoproduction off the proton target [7]. Besides the contributions of kaon trajectories and three hidden-charm pentaquarks, we revealed that the contact term plays a non-negligible role. For the time being, there are no experimental data on this process available. Thus, being limited by the broken SU(4) symmetry we adjusted the parameters of the model manually in order to provide at least qualitative predictions of the differential cross sections.

[1] D. Skoupil, P. Bydžovský, Phys. Rev. C 93, 025204 (2016).

[2] D. Skoupil, P. Bydžovský, Phys. Rev. C 97, 025202 (2018).

[3] P. Bydžovský, D. Skoupil, Phys. Rev. C 100, 035202 (2019).

[4] E. L. Lomon, Phys. Rev. C 66, 045501 (2002).

[5] H. Haberzettl et al., Phys. Rev. C 92, 055503 (2015).

[6] L. De Cruz et al., Phys. Rev. Lett. 108, 192002 (2012).

[7] D. Skoupil, Y. Yamaguchi, Phys. Rev. D 102, 074009 (2020).

Collaboration

Primary author: SKOUPIL, Dalibor (Nuclear Physics Institute CAS)

Presenter: SKOUPIL, Dalibor (Nuclear Physics Institute CAS)

Session Classification: Parallel Session C3