

Theoretical status of antikaon-nucleon interactions

Monday, 26 June 2023 09:00 (30 minutes)

In this talk, we discuss the recent theoretical studies of the strong interaction between an antikaon (\bar{K}) and a nucleon (N) [1,2]. The isospin $I = 0$ channel is of particular interest, as it exhibits an attractive interaction that can generate a quasi-bound state known as the $\Lambda(1405)$ resonance, located below the $\bar{K}N$ threshold. This suggests that the $\bar{K}N$ interaction may also produce quasi-bound states of kaonic nuclei [1]. Furthermore, the quasi-bound picture of the $\Lambda(1405)$ is related to the discussion of hadronic molecules in hadron spectroscopy.

Theoretical description of the $\Lambda(1405)$ in the coupled-channel meson-baryon scattering is developed with chiral SU(3) dynamics [2]. Based on the next-to-leading order (NLO) chiral SU(3) dynamics combined with the precise measurement of kaonic hydrogen by SIDDHARTA, it is shown that there are two resonances between the $\pi\Sigma$ and $\bar{K}N$ thresholds, $\Lambda(1405)$ and $\Lambda(1380)$. We review the current status of the theoretical studies of chiral SU(3) dynamics, including the recently performed NNLO analysis [3].

Femtoscopic study of the two-particle momentum correlation functions in high-energy collisions has become a new method to extract the hadron-hadron interactions. We present the study on the two-particle correlation function of a K^-p pair in high-energy collisions within the $\bar{K}N-\pi\Sigma-\pi\Lambda$ coupled-channels framework, which accurately account for all relevant effects, including the Coulomb potential and the threshold energy difference between K^-p and \bar{K}^0n [4]. Realistic $\bar{K}N-\pi\Sigma-\pi\Lambda$ potential based on NLO chiral SU(3) dynamics is used. We discuss the resulting K^-p correlation functions in comparison with the recent measurements by the ALICE collaboration under various collision conditions [4].

[1] T. Hyodo, W. Weise, arXiv:2202.06181 [nucl-th]

[2] U.G. Meißner, Symmetry 12, 981 (2020); M. Mai, Eur. Phys. J. ST 230 6, 1593 (2021); T. Hyodo, M Niiyama, Prog. Part. Nucl. Phys. 120, 103868 (2021).

[3] J.-X. Lu, L.S. Geng, M. Doering, M. Mai, Phys. Rev. Lett. 130, 071902 (2023).

[4] Y. Kamiya, T. Hyodo, K. Morita, A. Ohnishi and W. Weise, Phys. Rev. Lett. 124, 132501 (2020).

[5] S. Acharya et al. (ALICE collaboration), Phys. Rev. Lett. 124, 092301 (2020); S. Acharya et al. (ALICE collaboration), Phys. Lett. B 822, 136708 (2021); ALICE collaboration, arXiv:2205.15176 [nucl-ex]

Collaboration

Primary author: HYODO, Tetsuo (Tokyo Metropolitan University)

Presenter: HYODO, Tetsuo (Tokyo Metropolitan University)

Session Classification: Plenary session