Model selection in electromagnetic production of kaons

Dalibor Skoupil

(in collaboration with P. Bydžovský, A. Cieplý, D. Petrellis and D. Trnková)

Phys. Rev. C 93, 025204 (2016) Phys. Rev. C 97, 025202 (2018) Phys. Rev. C 104, 065202 (2021) Phys. Rev. C 107, 045206 (2023)

Department of Theoretical Physics Nuclear Physics Institute of the Czech Academy of Sciences, Řež, Czech Republic

17th International Workshop on Meson Physics, MESON 2023, Krakow, Poland June 26, 2023

Motivation for the work on kaon photo/electroproduction

- We aim at understanding the baryon spectrum and production dynamics of particles with strangeness at low energies.
- Constituent Quark Model predicts a lot more N* states than was observed in pion production experiments → "missing" resonance problem.
- Models for the description of elementary hyperon electroproduction are a suitable tool for hypernuclear physics calculations.
- New good-quality photoproduction data from BGOOD, LEPS, GRAAL, MAMI and (particularly) CLAS collaborations allow us to tune free parameters of the models.
- As the α_s increases with decreasing energy, we cannot use perturbative QCD at low energies \rightarrow need for introducing effective theories and models.

Introduction

Photoproduction process

 $p(p) + \gamma(k) \rightarrow K^+(p_K) + \Lambda(p_\Lambda)$

- Threshold: $E_{\gamma}^{lab} = 0.911 \text{ GeV}, W = 1.609 \text{ GeV}$
- In the lowest order, the reaction is described by the exchange of hadrons.
 - The 3rd nucleon-resonance region:

many resonant states and no dominant one in the kaon photoproduction \rightarrow need to assume a large number of nucleon resonances with mass $<2.5\,\text{GeV}$



 Resonance region: resonance contributions dominate (N*)

 Background: a plenty of nonresonant contributions (ρ, Κ, Λ; Κ* and Y*)

Isobar model

Single-channel approximation

 higher-order contributions (rescattering, FSI) included, to some extent, by means of effective values of the coupling constants

Use of effective hadron Lagrangian

- hadrons either in their ground or excited states
- amplitude constructed as a sum of tree-level Feynman diagrams
 - background and resonant part



· hadronic form factors account for the inner structure of hadrons

Free parameters (couplings, hff's cutoffs) adjusted to experimental data.

Satisfactory agreement with the data in the energy range $W = 1.6 - 2.5 \,\text{GeV}$.

Isobar model Calculation procedure

• Reaction amplitude: sum of *s*-, *t*-, and *u*-channel (non) Born amplitudes

$$\mathbb{M} = \sum_{x} \mathbb{M}_{x}$$
, where $x \equiv s, t, u, N^{*}, K^{*}, Y^{*}$

• Each contribution can be rewritten in a compact form

$$\mathbb{M}(\boldsymbol{\rho},\boldsymbol{\rho}_{\Lambda},\boldsymbol{k})=\bar{u}(\boldsymbol{\rho}_{\Lambda})\gamma_{5}\left(\sum_{j=1}^{6}\mathcal{A}_{j}(\boldsymbol{s},\boldsymbol{t},\boldsymbol{u})\mathcal{M}_{j}\right)\boldsymbol{u}(\boldsymbol{\rho}),$$

where A_j are scalar amplitudes and M_j are gauge-invariant operators, *i.e.* $k_{\mu}M_j^{\mu} = 0$,

$$\begin{split} \mathcal{M}_{1} &= \frac{1}{2} \left[\not k \not \in - \not \in \not k \right], & \mathcal{M}_{2} &= (p \cdot \varepsilon) - (k \cdot p) \frac{(k \cdot \varepsilon)}{k^{2}}, \\ \mathcal{M}_{3} &= (p_{\Lambda} \cdot \varepsilon) - (k \cdot p_{\Lambda}) \frac{(k \cdot \varepsilon)}{k^{2}}, & \mathcal{M}_{4} &= \not \in (k \cdot p) - \not k (p \cdot \varepsilon), \\ \mathcal{M}_{5} &= \not \in (k \cdot p_{\Lambda}) - \not k (p_{\Lambda} \cdot \varepsilon), & \mathcal{M}_{6} &= \not k (k \cdot \varepsilon) - \not \in k^{2}. \end{split}$$

Isobar model Calculation procedure

• CGLN amplitudes $f_i(k^2, s, t)$

$$\begin{split} \mathbb{M} &= \chi_{\Lambda}^{\dagger} \mathcal{F} \chi_{\rho}; \quad \mathcal{F} = f_{1}(\vec{\sigma} \cdot \vec{\varepsilon}) - if_{2}(\vec{\sigma} \cdot \hat{\rho}_{K})[\vec{\sigma} \cdot (\hat{k} \times \vec{\varepsilon})] + f_{3}(\vec{\sigma} \cdot \hat{k})(\hat{\rho}_{K} \cdot \vec{\varepsilon}) \\ &+ f_{4}(\vec{\sigma} \cdot \hat{\rho}_{K})(\hat{\rho}_{K} \cdot \vec{\varepsilon}) + f_{5}(\vec{\sigma} \cdot \hat{k})(\hat{k} \cdot \vec{\varepsilon}) + f_{6}(\vec{\sigma} \cdot \hat{\rho}_{K})(\hat{k} \cdot \vec{\varepsilon}) \end{split}$$

where e.g.

$$f_1 = N^*[-(W - m_p)\mathcal{A}_1 + (k \cdot p)\mathcal{A}_4 + (k \cdot p_\Lambda)\mathcal{A}_5 - k^2\mathcal{A}_6]$$

Response functions, e.g. transverse cross section

$$\begin{split} \frac{d\sigma}{d\Omega} &= \sigma_T = C \left\{ |f_1|^2 + |f_2|^2 - 2\operatorname{Re} f_1 f_2^* \cos \theta_K \\ &+ \sin^2 \theta_K \left[\frac{1}{2} (|f_3|^2 + |f_4|^2) + \operatorname{Re} \left(f_1 f_4^* + f_2 f_3^* + f_3 f_4^* \cos \theta_K \right) \right] \right\}, \end{split}$$

(for other response functions see Z. Phys. A 352 (1995) 327)

Isobar model Novel features of our isobar model

Exchanges of high-spin resonant states

non physical lower-spin components removed by appropriate choice of L_{int}

$$V^{\mu}_{S} \, \mathcal{P}^{(1/2)}_{ij,\,\mu
u} \, V^{
u}_{EM} = 0$$

Energy-dependent decay widths of nucleon resonances \rightarrow restoration of unitarity

$$\Gamma(\vec{q}) = \Gamma_{N^*} \frac{\sqrt{s}}{m_{N^*}} \sum_i x_i \left(\frac{|\vec{q}_i|}{|\vec{q}_i^{N^*}|} \right)^{2l+1} \frac{D(|\vec{q}_i|)}{D(|\vec{q}_i^{N^*}|)},$$

Extension from photoproduction to electroproduction

- · Phenomenological form factors in the electromagnetic vertex
- Longitudinal couplings of N^* 's to γ^* (crucial at small Q^2)

$$\begin{split} V^{EM}(N^*_{1/2} p\gamma) &= -i \frac{g_3^{EM}}{(m_R + m_p)^2} \Gamma_{\mp} \gamma_{\beta} \mathcal{F}^{\beta} \,, \\ V^{EM}_{\mu}(N^*_{3/2} p\gamma) &= -i \frac{g_3^{EM}}{m_R(m_R + m_p)^2} \gamma_5 \Gamma_{\mp} \left(\not a g_{\mu\beta} - q_{\beta} \gamma_{\mu} \right) \, \mathcal{F}^{\beta} \,, \\ V^{EM}_{\mu\nu}(N^*_{5/2} p\gamma) &= -i \frac{g_3^{EM}}{(2m_p)^5} \Gamma_{\mp} (q_{\alpha} q_{\beta} g_{\mu\nu} + q^2 g_{\alpha\mu} g_{\beta\nu} - q_{\alpha} q_{\nu} g_{\beta\mu} - q_{\beta} q_{\nu} g_{\alpha\mu}) \, p^{\alpha} \mathcal{F}^{\beta} \,. \end{split}$$

Fitting the data in the $K^+\Lambda$ channel

Minimization of χ^2 /n.d.f. with help of MINUIT library

Resonance selection

- s channel: spin-1/2, 3/2, and 5/2 N* with mass < 2.5 GeV;
- *t* channel: *K**(892), *K*₁(1272)
- *u* channel: *Y**(1/2) and *Y**(3/2)

Free parameters ($\approx 30 + 10$):

- SU(3)_f: $-4.4 \le g_{K\Lambda N}/\sqrt{4\pi} \le -3.0,$ $0.8 \le g_{K\Sigma N}/\sqrt{4\pi} \le 1.3$
- K*'s have vector and tensor couplings
- spin-1/2 resonance → 1 parameter; spin-3/2 and 5/2 resonance → 2 parameters
- 2 cut-off parameters for the hff
- 1 longitudinal coupling for each N*
- 2 cut-off parameters for the emff of K* and K₁

Experimental data

3383 $p(\gamma, K^+)\Lambda$ data

- cross section for W < 2.355 GeV (CLAS 2005 & 2010; LEPS, Adelseck-Saghai)
- hyperon polarisation for W < 2.225 GeV (CLAS 2010)
- beam asymmetry (LEPS)

171 $p(e, e'K^+)\Lambda$ data

σ_U, σ_T, σ_L, σ_{LT}, σ_K

Resulting models for the $K^+ \Lambda$ photo- and electroproduction



BS1 model ($\chi^2/n.d.f. = 1.64$)

- $S_{11}(1535), S_{11}(1650), F_{15}(1680), P_{13}(1720), F_{15}(1860), D_{13}(1875), F_{15}(2000);$
- K*(892), K₁(1272);
- Λ(1520), Λ(1800), Λ(1890), Σ(1660), Σ(1750), Σ(1940);
- multidipole form factor:

 $\Lambda_{bgr} = 1.88 \, \text{GeV}, \, \Lambda_{res} = 2.74 \, \text{GeV}$

BS3 model ($\chi^2/n.d.f. = 1.74$)

- $S_{11}(1535), S_{11}(1650), F_{15}(1680), P_{11}(1710), P_{13}(1720), F_{15}(1860), D_{13}(1875), P_{13}(1900), F_{15}(2000), D_{13}(2120);$
- K^{*}(892), K₁(1272);
- Λ(1405), Λ(1600), Λ(1890), Σ(1670);
- dipole form factor:

$$\Lambda_{bgr} = 1.24 \, \text{GeV}, \, \Lambda_{res} = 0.89 \, \text{GeV}$$

Transverse, σ_T , and longitudinal, σ_L , cross sections of $p(e, e'K^+) \wedge$



Extension from photo- to electroproduction

- BS1: naive extension by adding em. form factors only
- BS3: em. form factors and longitudinal couplings of N*'s to γ* added

 $\chi^{\rm 2}$ minimization and overfitting

Fitting procedure with MINUIT library: **minimizing the** χ^2

$$\chi^2 = \sum_{i=1}^{N} \frac{[d_i - p_i(c_1, \dots, c_n)]^2}{\sigma_{d_i}^2}$$

 (c_1, \ldots, c_n) - set of free parameters, (d_1, \ldots, d_N) - set of data points, p_i - theory, σ_{d_i} - error **Problem:** χ^2 minimization cannot prevent overfitting

Example: polynomial curve fitting

•
$$f(x, \mathbf{w}) = w_0 + w_1 x + w_2 x^2 + \dots + w_k x^k$$

• increasing order of polynomial *k* fits the data well...

...but gives only poor description of the function which generated them...

...and may fail to generalize to new data

 Occam's razor (law of parsimony): simpler models should be preferred



New fits of $K^+\Sigma^-$ channel Least Absolute Shrinkage and Selection Operator (LASSO)

Remedy to the overfitting issue: regularization

- introduce a penalty term to the $\chi^2 \rightarrow$ penalization of large parameter values
- penalized χ_P^2 : $\chi_P^2 = \chi^2 + P(\lambda)$
- penalty term: $P(\lambda) = \lambda^4 \sum_{i=1}^{N_{res}} |g_i|$
 - λ regularization parameter, g_i resonances' couplings
- LASSO forces some of the parameters to zero
 → selection of a subset of the fit parameters
- λ controls the strength of the penalty and thus the complexity of the model
 → higher powers of λ allow fine sampling of the region of small λ

Information criteria:

- Akaike information criterion AIC = $2n_i + \chi_P^2$
- Corrected Akaike information criterion AICc = AIC + $\frac{2n_i(n_i+1)}{N-n_i-1}$
- Bayesian information criterion BIC = $n_i \ln(N) + \chi_P^2$

 n_i - no. of parameters corresponding to λ_i

N - number of data points



Applying the information criteria - forward selection

- 1 start with the full model: parameters initialized within $\langle -1; +1 \rangle$; use λ_{max}
- 2 perform LASSO χ^2_P minimization and compute IC
- ${f 3}$ in each run reduce λ and run LASSO with the values of the previous run as starting values
- 4 repeat until λ_{\min} is reached

Optimal λ occurs at the minimum of the IC.

Fitting procedure

- resonance selection: motivation from previous analysis of $K^+\Lambda$ channel
- non resonant part: Born terms and exchanges of K* and K₁ and Σ*'s
- resonant part: exchanges of N^* 's and Δ^* 's in the s channel
- around 600 data utilized to fit \leq 25 parameters
- result with the smallest $\chi^2/ndf = 2.3 \rightarrow fit M$ (25 parameters, 14 resonances)
- LASSO applied at fit M: χ^2_P /ndf = 3.4 \rightarrow fit L (17 parameters, 9 resonances)

Characteristics of models

- only one Δ resonance introduced
- no hyperon resonances needed for reliable data description
- results in very good agreement with the cross-section and beam-asymmetry data
- fit L is very economical

Differential cross section in dependence on the photon lab energy



Differential cross section in dependence on the photon lab energy - fit L w/o individual resonances



Notation: N7: N(1720)3/2⁺, M4: N(2060)5/2⁻

Dalibor Skoupil

Beam asymmetry in dependence on the kaon center-of-mass angle - fit L w/o individual resonances



LASSO in the $K^+\Lambda$ channel

Selecting a subset of resonances (results very preliminary!)





Resonances in BS1 and BS1L:

- $S_{11}(1535), S_{11}(1650), F_{15}(1680), P_{13}(1720), F_{15}(1860), D_{13}(1875), F_{15}(2000);$
- K*(892), K₁(1272);
- Λ(1520), Λ(1800), Λ(1890), Σ(1660), Σ(1750), Σ(1940)

Model	no. of resonances	no. of parameters	$\chi^2/{\rm n.d.f.}$
BS1	16	31	1.6
BS1L	9	20	3.6

Refitting the model's parameters in the $K^+\Lambda$ channel

Ridge regression and cross validation for suppressing hyperon couplings

Why refit?

- include recent measurements of polarization observables (PR C 93, (2016) 065201)
- need to investigate more the role of hyperon resonances in KY photoproduction
- large values of hyperon couplings: ridge regression to suppress them during the fitting procedure

Ridge regularization

- penalized χ_P^2 : $\chi_P^2 = \chi^2 + \lambda^4 \sum_{i=1}^{n_{\Lambda}} g_i^2$, ($n_{\Lambda} = no. \text{ of } Y \text{ couplings}$)
- parameter values reduced but they are not reduced to zero

Cross validation







Model selection in elmag production of kaons

"One standard-error rule" for selecting appropriate model

Errors on the validation set help us determine the optimal value of $\boldsymbol{\lambda}$

• average of validation errors over all k runs

$$\overline{CV}(\lambda) = \frac{1}{k} \sum_{j=1}^{k} CV_j(\lambda)$$

• optimal
$$\lambda^*$$
 is then:

$$\lambda^* = \operatorname*{argmin}_{\lambda \in \{\lambda_{\min}, \dots, \lambda_{\max}\}} \overline{CV}(\lambda)$$

"One standard-error rule":

the most parsimonious model within one standard error from the minimum of the validation error should be chosen

new optimal λ̃ is such that

$$\overline{CV}(\tilde{\lambda}) = \overline{CV}(\lambda^*) + SE(\lambda^*)$$



Relative percentage reduction of the resonance couplings



BS2

BS2r

Tag	Resonance	g_1	g_2		
L1	$\Lambda(1405) \ 1/2$	9.67	-	2.624	
S1	$\Sigma(1660) \ 1/2^+$	-8.09	-	 -5.925	
L4	$\Lambda(1800) \ 1/2^{-}$	-11.55	-	-1.409	
S4	$\Sigma(1940) \; 3/2^-$	-0.86	0.18	-0.685	0.079

- g_i values from the unregularized fitting
- \tilde{g}_i values after performing Ridge regularization

$K^+\Lambda$ channel: beam asymmetry Σ



$K^+\Lambda$ channel: target asymmetry T



Summary

New version of isobar model for the $K^+\Lambda$ channel

available for calculations online at:

```
http://www.ujf.cas.cz/en/departments/
department-of-theoretical-physics/isobar-model.html
```

Description extended from the $K^+\Lambda$ channel to the $K^+\Sigma^-$ channel.

Regularization methods introduced as a remedy for overfitting and as model selection tools.

Outlook

- testing the models in the DWIA calculations for hypernucleus production
- performing an analysis of Σ photoproduction channels
- extending the analysis of electroproduction beyond $Q^2 = 1 \text{ GeV}^2$
- studying the production of Ξ hypernuclei
- etc., etc....

Thank you for your attention!